

STRICT DEAD END ELEMENTS IN FREE SOLUBLE GROUPS

VICTOR GUBA

ABSTRACT. Let G be a group generated by a finite set A . An element $g \in G$ is a strict dead end of depth k (with respect to A) if $|g| > |ga_1| > |ga_1a_2| > \cdots > |ga_1a_2 \cdots a_k|$ for any $a_1, a_2, \dots, a_k \in A^{\pm 1}$ such that the word $a_1a_2 \cdots a_k$ is freely irreducible. (Here $|g|$ is the distance from g to the identity in the Cayley graph of G .) We show that in finitely generated free soluble groups of degree $d \geq 2$ there exist strict dead elements of depth $k = k(d)$, which grows exponentially with respect to d .

Let G be a group generated by a finite set A . By $\mathcal{C} = \mathcal{C}(G, A)$ we denote the *right Cayley graph* of G with respect to A . The set of vertices of this graph is G , the set of edges is $G \times A^{\pm 1}$. Every edge $e = (g, a)$, where $g \in G$, $a \in A^{\pm 1}$, starts in g and ends in ga . This edge is labelled by a . An inverse edge is $e^{-1} = (ga, a^{-1})$. For every $g \in G$ let $|g|$ denote the distance in $\mathcal{C}$ from the vertex g to the identity.

An element g is a *dead end* in $\mathcal{C}$ whenever $|g| > |ga|$ for all $a \in A^{\pm 1}$. This concept was introduced by Bogopolski in [3]. One of the easiest examples is the following. Let us consider the group $G = \mathbb{Z} \times \mathbb{Z}_2$, where a, b generate the factors. It is easy to see that the Cayley graph of G in generators a, b has no dead end vertices. However, if the generating set is $A = \{a, ab\}$, then b becomes a dead end element. This example shows that being a dead end element strongly depends on the generating set. However, if the generating set is clear, we will often say that an element of G is (or is not) a dead end.

A less trivial example of dead end elements comes from the R. Thompson group F . Recall that F is the group presented by

$$\langle x_0, x_1, x_2, \dots \mid x_j x_i = x_i x_{j+1} \ (j > i) \rangle.$$

This group can be generated by x_0, x_1 . In these generators, the group has infinitely many dead end elements. Say, $g = x_0^2 x_1 x_6 x_3^{-1} x_0^{-2}$ will be a dead end element of length 11. One can see details in [11]. (There are no dead end elements in F of length less than 11.) There are descriptions of all dead end elements in F given independently in [2]

2000 *Mathematics Subject Classification.* Primary 20F32; Secondary 05C25.

Key words and phrases. Cayley graphs; dead end elements; free soluble groups.

This research is partially supported by the RFFI grant 05-01-00895 and the INTAS grant 99-1224.

and [5]. They are based on the representations of F in terms of (pairs of) rooted binary forests and trees, respectively.

One can try to generalize the concept of a dead end element in different ways. Say, one can ask if there are elements in F with the following property: $|g| > |ga| > |gab|$ for all $a, b \in \{x_0^{\pm 1}, x_1^{\pm 1}\}$ such that a, b are not inverses of each other. It can be shown that such elements do not exist [5].

One of the generalizations has been considered in [6], see also [4]. Let G be a group and let A be its finite generating set. An element $g \in G$ is a *dead end of depth k* in the Cayley graph $\mathcal{C}(G, A)$ whenever the ball of radius k around g is contained in the ball of radius $|g|$ around the identity. In other words, $|g| \geq |ga_1|, |ga_1a_2|, |ga_1a_2 \cdots a_k|$ for all $a_1, a_2, \dots, a_k \in A^{\pm 1}$. Hence each dead end element is a dead end of depth 2. It follows from [5] that F has no dead ends of depth ≥ 3 (the set of generators is $\{x_0, x_1\}$ as above).

One can ask whether there are groups whose Cayley graphs contain dead end elements of arbitrary depth. A positive answer is given in [6], where the group is the *lamplighter group*, which is the restricted wreath product $\mathbb{Z}_2 \wr \mathbb{Z}$ with natural set of generators. Finitely presented examples of that kind are considered in [4].

One more natural generalization comes as follows. Let us say that an element $g \in G$ is a *strict dead end of depth k* in the Cayley graph $\mathcal{C}(G, A)$ whenever $|g| > |ga_1| > |ga_1a_2| > \cdots > |ga_1a_2 \cdots a_k|$ for all $a_1, a_2, \dots, a_k \in A^{\pm 1}$ such that the word $a_1a_2 \cdots a_k$ is freely irreducible. This means that if we travel in $\mathcal{C}$ from g along a path of length k with freely irreducible label, then we become closer and closer to the identity each time we go through an edge. Obviously, each dead end element is exactly a strict dead end element of depth 1.

Unlike the case of (non-strict) dead end elements, it is obvious that any group can have strict dead elements of bounded depth only. Indeed, a group freely generated by A has no dead ends at all. If there is a nontrivial group relation between elements of A , then let us take the shortest one. If ρ denotes its length, then there are no strict dead end elements of depth $k > \lfloor \rho/2 \rfloor$. The goal of this article is, given a k , to construct a group that has strict dead elements of depth k . Moreover, we find such examples among free soluble groups. These groups are not finitely presented if the degree of solvability is greater than or equal to 2 (see [16]). However, an easy observation shows that the needed examples can be found in the class of finitely presented groups and even finite groups.

We need to recall some facts from the Stallings theory of group ends. Let G be a finitely generated group. Let us consider its Cayley graph $\mathcal{C} = \mathcal{C}(G, A)$ and take the ball B_n of radius n around the identity. The graph $\mathcal{C} - B_n$ (we delete all vertices at distance $\leq n$ from the identity

and all edges coming out of them) may consist of several connected components. If n approaches infinity, then the number of infinite connected components of $\mathcal{C} - B_n$ approaches some value that should be equal to 0, 1, 2, or infinity. This is the number of *ends* of the group G . It is independent on the choice of the finite generating set for G . It is obvious that G has zero ends if and only if it is finite. If G has two ends, then it is a finite extension of $\mathbb{Z}$. If G has infinitely many ends, then G is a nontrivial amalgamated product or a nontrivial HNN-extension with finite amalgamated (associated) subgroup(s) [17]. Therefore, in many cases we can conclude that a group has only one end.

Now we need to recall some facts from [8] that will play an important rôle for us. For some of them, we recall the idea of the proof for reader's convenience.

Let $\mathbb{F}_m$ be a free group with m generators and let R be its normal subgroup. We fix the standard generating set A for $\mathbb{F}_m$. This will also denote the generating sets for quotient groups $\mathbb{F}_m/R$ and $\mathbb{F}_m/R'$, where $R' = [R, R]$ is the commutator subgroup of R .

By $\text{Sol}(m, d)$ we denote the *free soluble* group of rank m and *degree of solvability* d . This is the quotient $\mathbb{F}_m/\mathbb{F}_m^{(d)}$, where $H^{(d)}$ denotes the d th *derived subgroup* of H . This is defined inductively: $H^{(0)} = H$ and $H^{(n+1)} = [H^{(n)}, H^{(n)}]$ for all $n \geq 0$. To work with free soluble groups, one needs to describe the connection between groups $\mathbb{F}_m/R$ and $\mathbb{F}_m/R'$ in its general form.

There is a natural way to describe the solution to the word problem in $\mathbb{F}_m/R'$ based on some properties of the Cayley graph of $\mathbb{F}_m/R$. Moreover, the paper [8] contains an algorithm how to find the length of a given element in $\mathbb{F}_m/R'$ with respect to A . We need to introduce some terminology.

Let Γ be a graph in the sense of Serre [15] (that is, each edge has an inverse). A *flow* on Γ is a function μ from the set of edges $E = E(\Gamma)$ to $\mathbb{R}$ such that $\mu(e^{-1}) = -\mu(e)$ for all $e \in E$. We say that $\mu(e)$ is a flow *through* the edge e . If Γ is locally finite, then we can define an *inflow* $\text{infl}_\mu(v)$ into a vertex v . This is the sum of flows through all edges e that have v as its terminal vertex.

Given a flow μ on a graph, we will say that its *support* $\text{supp}(\mu)$ is a subgraph in Γ that is the union of all the edges e satisfying $\mu(e) \neq 0$ together with their endpoints. Thus μ has finite support if and only if $\mu(e) = 0$ for all but finitely many $e \in E$. For an edge $e \in E$, we denote by χ_e the *characteristic flow* of e on Γ . Namely, $\chi_e(e) = 1$, $\chi_e(e^{-1}) = -1$, $\chi_e(x) = 0$ otherwise. Obviously, the set of flows on Γ forms a vector space. If we fix an orientation on Γ (that is, if we choose a subset E_+ in E such that E_+ contains exactly one of the edges e, e^{-1} for all $e \in E$), then the set of flows $\chi_e, e \in E_+$ forms a basis of the subspace of flows with finite support.

A flow on Γ is called *balanced* whenever the inflow into any vertex is zero. A flow μ on Γ is *semi-balanced* whenever there exist two vertices v_- , v_+ such that $\text{infl}_\mu(v_-) = -1$, $\text{infl}_\mu(v_+) = 1$ and $\text{infl}_\mu(x) = 0$ for all $x \neq v_-, v_+$. Every path p in Γ induces a flow χ_p on Γ in the following way: if $p = e_1 \cdots e_n$ is a product of edges, then $\chi_p = \chi_{e_1} + \cdots + \chi_{e_n}$ by definition. Such a flow is always balanced or semi-balanced.

Given a flow μ , an edge is called a *P-edge* whenever $\mu(e) > 0$. If $\text{supp}(\mu)$ is finite, then the sum of flows through all the *P-edges* is called the *weight* of μ .

Lemma 1. *Let μ a nonzero integer-valued flow on a graph Γ . Suppose that μ is balanced and let $\text{supp}(\mu)$ be finite and connected. Then for any vertex v of $\text{supp}(\mu)$, there exists a loop p at v in Γ such that p is a product of *P-edges* and every edge $e \in E$ occurs in this product exactly $\mu(e)$ times.*

Proof. Let N be the weight of μ . We proceed by induction on N . Let e_1 be any *P-edge*. For every $i \geq 1$, let v_i denote the terminal vertex of e_i . Since the inflow into v_i is zero, there exists a *P-edge* e_{i+1} starting at v_i . Since the support of μ is finite, some edges of the form e_i ($i \geq 1$) will repeat. Hence we can find a closed path of the form $q = e_{i+1} \cdots e_{i+k}$, $k \geq 1$ that consists of different *P-edges*.

Let $\mu' = \mu - \chi_q$. Clearly, μ' is also balanced and has a finite support. The weight of μ' equals $N - k < N$. If μ' is zero, then we are done. Otherwise $\text{supp}(\mu')$ is a union of its connected components $\Sigma_1, \dots, \Sigma_s$ ($s \geq 1$). We can apply the inductive assumption to the restriction of μ' on each of the components.

It is easy to see that each Σ_j ($1 \leq j \leq s$) has a common vertex v_j with the path q . These vertices are different for different values of j . Changing the order of the components, we can assume that $q = q_0 q_1 \cdots q_s$, where v_j is the initial point of q_j ($1 \leq j \leq s$). By the inductive assumption, there is a loop p_j at v_j that contains each *P-edge* e of Σ_j exactly $\mu'(e)$ times ($1 \leq j \leq s$). Now the path $p = q_0 p_1 q_1 \cdots p_s q_s$ is the loop we wanted to find. $\square$

Let G be a group generated by a finite set A . For every group word w over A there exists a unique path $p = p(w)$ in the Cayley graph $\mathcal{C} = \mathcal{C}(G, A)$ starting at the identity and labelled by w . So we can assign to w the flow $\mu_{p(w)}$ on $\mathcal{C}$. This flow is called the *flow induced by the word w* . It is obvious that if two words are equal in the free group, then they induce the same flow. It is also obvious that the flow induced by a word has finite support. It is balanced whenever $p = p(w)$ is closed (that is, w equals 1 in G) and semi-balanced whenever $p = p(w)$ is not closed (in this case $v_- = 1$, $v_+ = g$, where $g \in G$ is the element represented by w). It is easy to prove the following fact.

Lemma 2. *A flow μ on $\mathcal{C} = \mathcal{C}(G, A)$ is induced by a word if and only if the following conditions hold:*

- μ is integer-valued,
- μ has finite support,
- μ is balanced or it is semi-balanced with $v_- = 1$.

Proof. The “only if” part is obvious. Now assume that a flow μ satisfies all the three conditions. First of all, let μ be balanced. If $\text{supp}(\mu)$ is connected, we find a path q from the identity to a point v in $\text{supp}(\mu)$ and a loop p from Lemma 1. Then the label of qpq^{-1} will induce the desired flow. If $\text{supp}(\mu)$ is not connected, then we take a product of words that induce the flow for each of the connected components.

In the semi-balanced case, let us assume that μ has the property $\text{infl}_\mu(1) = -1$, $\text{infl}_\mu(g) = 1$ with zero inflow into all the other vertices. Let us consider any path q labelled by a word Q that connects g and 1. Let us add the flow χ_q to μ . This flow will be balanced and thus induced by a word w . Clearly, the word wQ^{-1} will induce μ . $\square$

The following fact has been proved in [8].

Lemma 3. *Let R be a normal subgroup in the free group $\mathbb{F}_m$. Two words w_1, w_2 are equal modulo the commutator subgroup R' if and only if they induce the same flow on the Cayley graph of $\mathbb{F}_m/R$ with respect to the natural set of generators.*

For the sake of completeness, let us recall the idea of the proof. A word w belongs to R if and only if the path $p = p(w)$ is a loop in the Cayley graph $\mathcal{C}$ of $\mathbb{F}_m/R$. Hence any word of the form $[r_1, r_2] = r_1^{-1}r_2^{-1}r_1r_2$, where $r_1, r_2 \in R$, will induce a zero flow on $\mathcal{C}$. The same holds for a product of these commutators, which implies the “only if” part. To prove the converse, it suffices to choose a spanning subtree T in $\mathcal{C}$. For each edge e , one can define a loop $\bar{e} = peq^{-1}$ at the identity, where p, q are geodesic paths in T that connect 1 with the initial and the terminal vertex of e , respectively. If w_1, w_2 induce the same flow on $\mathcal{C}$, then the paths $p(w_1), p(w_2)$ have the same endpoints. Hence $s = p(w_1)p(w_2)^{-1}$ will be the loop at the identity. If $s = e_1 \cdots e_n$ is the product of edges, then s is freely homotopic to the path $\bar{e}_1 \cdots \bar{e}_n$. Since the flow χ_s is zero, the number of occurrences of e in s is the same as the number of occurrences of e^{-1} , for any e . Taking into account that $\bar{e}$ represents an element of R and the fact that $\overline{e^{-1}} = \bar{e}^{-1}$, we conclude that the label of s equals 1 modulo R' (we do permutations in the product $\bar{e}_1 \cdots \bar{e}_n$ cancelling each $\bar{e}$ with its inverse).

The paper [8] also has a description of lengths of all elements in $\mathbb{F}_m/R'$ and their minimal word representatives. This means that, given a word w , we can find a word v that equals w modulo R' and has the shortest possible length. (This word is not unique in general.) The length of the shortest representative of g is denoted by $|g|$ and it is called the *length* of the element $g \in \mathbb{F}_m/R'$. In fact, the algorithm to find the length of an element given by a word w is polynomial on $|w|$.

(One can compare this situation with the one described in [14], where it is shown that the problem to find the length of an element of the restricted wreath product $\mathbb{Z} \wr \mathbb{Z}^2$ in its natural generators turns out to be NP-complete.)

We need an explicit description of lengths of elements in $\mathbb{F}_m/R'$ of some particular form.

Lemma 2 shows us that any balanced flow with finite support on the Cayley graph of $\mathbb{F}_m/R$ is induced by a word w . It follows from Lemma 3 that w is defined uniquely modulo R' . Therefore, we can say that any flow with the above properties defines canonically an element of $\mathbb{F}_m/R'$. (In fact, all these elements form the subgroup R/R' .)


The following statement is a partial case of [8, Theorem 2].

Lemma 4. *Let μ be a nonzero balanced flow with finite support on the Cayley graph of $\mathbb{F}_m/R$. Let $g \in \mathbb{F}_m/R'$ be defined by this flow. Suppose that $\text{supp}(\mu)$ is connected. Let d be the distance from the identity to $\text{supp}(\mu)$ in the above Cayley graph. Then the length of the shortest word that represents g in $\mathbb{F}_m/R'$ is given by the formula $|g| = N + 2d$, where N is the weight of μ .*

Proof. Suppose that a word w induces the flow from the statement. Let $p = p(w)$ be the path labelled by w starting at the identity in the Cayley graph $\mathcal{C}$ of the group $\mathbb{F}_m/R$. Let q be the shortest subpath in p that contains all edges with nonzero flow. Obviously, $|q| \geq N$. If $p = p'qp''$, then it is clear that $|p'|, |p''| \geq d$. This proves that $|w| \geq N + 2d$.

Now let p_0 be a path of length d that connects the identity with some vertex v from $\text{supp}(\mu)$. We know from Lemma 1 that there is a loop p at v that consists of N edges such that each P -edge e occurs in p exactly $\mu(e)$ times. The path $p_0pp_0^{-1}$ has length $N + 2d$ and its label represents g . Therefore, $|g| = N + 2d$. $\square$

Now we are ready to present an example of a strict dead end element of depth 2 in a free metabelian group. For simplicity, we consider the group with 2 generators. Let μ be a balanced flow given by the “thick” curve on the picture below. The curve or one of its appropriate cyclic shifts read in the clockwise direction will be denoted by p .



It is easy to see that the flow μ determines an element $g \in G = \mathbb{F}_2/\mathbb{F}_2''$, where $g = b^2(ab^{-1})^2(b^{-1}a^{-1})^2(a^{-1}b)^2(ba)^2b^{-2}$. According to Lemma 4, the length of g in G equals 20. In fact, there are many ways to express g by a word of minimal length. One can take any path q that connects the identity to a point on the curve p , then go around the curve and return back to the identity by the path q^{-1} . It is easy to see that the element gx has length 19 for any $x \in \{a^{\pm 1}, b^{\pm 1}\}$. Indeed, one can choose q in such a way that the first edge e of q has label x . If $q = eq'$, then the path $qp(q')^{-1}$ representing gx has length 19. Therefore, the length of gx equals 19 (it cannot decrease by more than 1).

Now let xy be any freely irreducible word of length 2 over the alphabet $\{a^{\pm 1}, b^{\pm 1}\}$. Clearly, the path q of length 2 labelled by xy starting at the identity will end on the curve p . Then gxy is represented by the path qp of length 18, which is also the length of gxy in G .

By definition, $g \in G$ is a strict dead end element of depth 2. It is easy to see that g is not a strict dead end of depth 3 since $gaba^{-1}$ has length 19.

A group word in the alphabet $\{x_1, \dots, x_m\}$ is called *positive* whenever it does not contain letters of the form x_i^{-1} ($1 \leq i \leq m$). Here is one of the main results.

Theorem 1. *Let R be a normal subgroup in the free group $\mathbb{F}_m$ of rank $m > 1$. Assume that the following three conditions hold:*

- 1) *the group $\mathbb{F}_m/R$ has exactly one end,*
- 2) *there are no nonempty positive words in R ,*
- 3) *any nonempty cyclically reduced word in R has length at least ρ .*

Then the group $G = \mathbb{F}_m/R'$ has strict dead end elements of depth $\lceil \rho/2 \rceil - 1$.

Proof. Let $k = \lceil \rho/2 \rceil - 1$. It follows from the third condition that the ball B_k of radius k around the identity is a tree. Let $\mathcal{C}$ be the Cayley graph of $\mathbb{F}_m/R$ and let $\mathcal{C}'$ be the subgraph $\mathcal{C} - B_{k-1}$. We claim that $\mathcal{C}'$ is connected.

Assume the contrary. Then $\mathcal{C}'$ has more than one connected component. Only one of them is infinite because of the first condition. So there is a finite component Γ in $\mathcal{C}'$. Each vertex in $\mathcal{C}'$ has degree at least $2m - 1 > m$ since $m \geq 2$. Let us take any edge e_1 in Γ with positive label and let v_1 be its terminal vertex. There exists at least one edge e_2 with positive label that comes out of v_1 . (Otherwise v_1 has degree at most m .) Denoting the terminal vertex of e_2 by v_2 , we see that there is an edge e_3 that comes out of v_2 and has a positive label, and so on. Since Γ is finite, we will have repetition of vertices on some step and so there will be a positive nontrivial relation in R , which contradicts the second condition.

Now we want to prove a slightly stronger fact. Suppose that e is an edge that belongs to $\mathcal{C}'$. We want to show that the graph $\mathcal{C}' - e$ (we

remove the edges $e^{\pm 1}$) is still connected. Assume the contrary. Suppose that the endpoints of e belong to different connected components of $\mathcal{C}' - e$. One of these components must be finite. Let v_0 be the endpoint of e that belongs to this finite component Γ . All vertices in Γ still have degree at least $2m - 1$ in this component except for, possibly, v_0 , which has degree at least $2m - 2$. Let e_1 be an edge that comes out of v_0 . If this edge has a positive label, then we can repeat the arguments from the previous paragraph. If the edge e_1 has a negative label, then we repeat the same arguments replacing “positive” by “negative” everywhere and get a contradiction.

The fact that $\mathcal{C}' - e$ is always connected means that each edge e in $\mathcal{C}'$ belongs to a simple loop contained in $\mathcal{C}'$. We want to show that $\mathcal{C}'$ contains a finite subgraph with certain properties. First of all, we connect each pair of vertices at a distance k from the identity by a path in $\mathcal{C}'$. This gives a finite subgraph in $\mathcal{C}'$. Now, for each edge of this subgraph, we choose a simple loop in $\mathcal{C}'$ that contains this edge. We add all these loops to the subgraph and get a new finite connected subgraph Σ . By the construction, each edge of Σ is contained in a simple loop that is contained in Σ .

Let $e_1, e_2, \dots, e_t$ be all positively labelled edges of Σ and let $q_1, q_2, \dots, q_t$ be simple loops that contain these edges, respectively (we allow some of the loops coincide). For every $1 \leq i \leq t$, let us consider the flow $2^i \chi_{q_i}$ and let μ be the sum of these flows. Clearly, μ is balanced and $\mu(e_i) \neq 0$ for all $1 \leq i \leq t$. Indeed, for any e_i let j be the smallest number such that $e_i^{\pm 1}$ occurs in q_j . Clearly, $j \leq i$ and $\mu(e_i)$ equals 2^j modulo 2^{j+1} .

The identity element is at a distance k from Σ . If N is the weight of μ , then the element g in $\mathbb{F}_m/R'$ represented by μ has length $N + 2k$ in this group according to Lemma 4. We claim that g is a strict dead end element of depth at least k in $\mathbb{F}_m/R'$.

Indeed, let $a_1 \cdots a_k$ be any freely irreducible word in the generators of G . We want to prove that $|g| > |ga_1| > |ga_1a_2| > \cdots > |ga_1a_2 \cdots a_k|$. We show that $|ga_1 \cdots a_i| = |g| - i$ for all $0 \leq i \leq k$. Obviously, $|ga_1 \cdots a_{i-1}a_i| \geq |g| - i$. On the other hand, let q be the path of length k that connects 1 to a vertex v in Σ such that $q = q'q''$ and the label of q' is $a_1a_2 \cdots a_i$. By Lemma 1, there exists a loop p at v of length N that induces μ . Then the label of the path $qp(q'')^{-1}$ represents $ga_1 \cdots a_i$. The length of this path is $|g| - |q'| = |g| - i$. This implies what we wanted to prove. $\square$


Let us estimate the length ρ_d of a shortest nontrivial relation in $\text{Sol}(m, d)$, where $m > 1$. It is not hard to see that $\rho_1 = 4$, $\rho_2 = 14$. In general, we have the following rough estimate.

Lemma 5. *Let ρ_d be the length of the shortest nontrivial relation in the free soluble group $\text{Sol}(m, d)$, where $m > 1$, $d \geq 1$. Then $\rho_{d+1} \geq 3\rho_d$ for all $d \geq 1$.*

Proof. Let $R = \mathbb{F}_m^{(d)}$. Suppose that w is a shortest nontrivial word that represents 1 in $\mathbb{F}_m/R$. By Lemma 3, w induces a zero flow on the Cayley graph $\mathcal{C}$ of $\mathbb{F}_m/R$. Let $p = p(w)$ be the path in $\mathcal{C}$ labelled by w starting at 1. Clearly, p cannot have (cyclic) subpaths of the form ee^{-1} , where e is an edge. (Otherwise w is not a shortest nontrivial relation.)

Let Γ be a subgraph in $\mathcal{C}$ formed by all the edges that occur in p . It is obvious that Γ is connected; it is also clear that Γ has no vertices of degree 1.

One can observe that Γ is not a circle, that is, it cannot have rank 1 (a rank of a connected graph is the rank of its fundamental group). Hence the rank of Γ is at least 2. It is well known that each graph of rank greater than 1 has a (geometric) subgraph of one of the following forms (cf. [9]):



We see that a) can be considered as a partial case of b) with r as the empty path. Since p goes at least once through each edge of Γ in both directions, one can estimate the length of p as $|p| \geq 2(|s| + |t| + |r|)$ in b) and c). Since every nontrivial relation in $\mathbb{F}_m/R$ has length at least ρ_d , we can conclude that $|s|, |t| \geq \rho_d$ in the case b) and so $|p| \geq 4\rho_d$. In the case c), we have loops sr , $r^{-1}t$, and st , each of length at least ρ_d . Hence $|p| \geq 2(|s| + |t| + |q|) = (|s| + |q|) + (|q| + |t|) + (|s| + |t|) \geq 3\rho_d$. Therefore, we always have $\rho_{d+1} \geq 3\rho_d$. $\square$

Lemma 5 immediately implies that $\rho_d \geq 4 \cdot 3^{d-1}$ for the group $\text{Sol}(m, d)$, where $m > 1$, $d \geq 1$.

Now we can apply Theorem 1 to extract a corollary for the case of free soluble groups.

Theorem 2. *If $m > 1$, $d \geq 2$, then the free soluble group $\text{Sol}(m, d)$ has strict dead end elements of depth at least $2 \cdot 3^{d-2} - 1$.*

Proof. Free soluble groups have only one end in the sense of Stallings. Indeed, they are not virtually cyclic and they cannot be presented as a nontrivial amalgamated product or an HNN-extension. Let $R = \mathbb{F}_m^{(d-1)}$. Every nontrivial word in the commutator subgroup of $\mathbb{F}_m$ has zero exponent sum on each generator. Hence the only positive word in $R \subseteq \mathbb{F}_m'$ is the empty word. It follows from Lemma 5 that the length of the shortest nontrivial relation in R is $\rho_{d-1} \geq 4 \cdot 3^{d-2}$. Now we apply

Theorem 1 and conclude that the group $\text{Sol}(m, d) = \mathbb{F}_m/R'$ has strict dead end elements of depth at least $\lceil \rho_{d-1}/2 \rceil - 1 \geq 2 \cdot 3^{d-2} - 1$. $\square$

Notice that the groups $\text{Sol}(m, d)$ are not finitely presented whenever $m > 1$, $d \geq 2$ by a result of [16]. However, they are residually finite [10]. Therefore, for any $k \geq 1$ there exist finitely presented and even finite groups with strict dead end elements of depth k .

Remark. The solution to the word problem in free soluble groups gives rise to a question about growth functions for these groups. Recall that if $\mathcal{C} = \mathcal{C}(G, A)$, then the *growth function* of G with respect to A is the sequence $b_n = \text{Card}(B_n)$, where B_n denotes the ball of radius n in $\mathcal{C}$. The limit $\beta = \lim_{n \rightarrow \infty} \sqrt[n]{b_n}$ always exists; it is called the *growth rate* of G (with respect to A).

Even in the case of $G = \text{Sol}(2, 2)$, a 2-generated free metabelian group, neither the growth function nor the growth rate is known. A lower bound for the growth function of G is the number of self-avoiding walks (SAW) of length n in the square lattice starting at the identity (different self-avoiding paths represent different elements of G by Lemma 3). The exact value of the growth rate of SAW is not known although more than 10 of its decimal digits were found [12]. So the growth rate of G satisfies inequality $\beta \geq 2.63815853034$.

Notice that for some metabelian 2-generated groups both the growth rate and the growth function are known. Say, Parry [14] found them for the restricted wreath product $\mathbb{Z} \wr \mathbb{Z}$ and for some other groups. For $\mathbb{Z} \wr \mathbb{Z}$, the growth function is rational and the growth rate equals $\sqrt{2} + 1$. In [7] it is shown that the growth functions of Baumslag – Solitar groups $BS(1, n)$ are rational; their growth rates do not exceed the above value $\sqrt{2} + 1$.

In a recent paper [1], the authors prove that the growth rate of the free soluble group $\text{Sol}(m, d)$ approaches $2m - 1$ as $d \rightarrow \infty$. This shows that the growth rate of amenable groups with m generators can be arbitrarily close to the maximum value $2m - 1$.

REFERENCES

- [1] G. N. Arzhantseva, V. S. Guba, L. Guyot. Growth rates of amenable groups. J. Group Theory 8, no. 3 (2005), 389–394.
- [2] James M. Belk and Kenneth S. Brown. Forest diagrams for elements of Thompson’s group F .
<http://www.arxiv.org/abs/math.GR/0305412>
- [3] O. V. Bogopolski. Infinite commensurable hyperbolic groups are bi-Lipschitz equivalent. Algebra and Logic 36, no. 3 (1997), 155–163.
- [4] Sean Cleary and Tim R. Riley. A finitely presented group with unbounded dead-end depth.
<http://www.arxiv.org/abs/math.GR/0406443>
- [5] Sean Cleary and Jennifer Taback. Combinatorial properties of Thompson’s group F . Trans. Amer. Math. Soc. (to appear).

- [6] Sean Cleary and Jennifer Taback. Dead end words in lamplighter groups and other wreath products.
<http://www.arxiv.org/abs/math.GR/0309344>
- [7] D. J. Collins, M. Edjvet and C. P. Gill. Growth series for the group $\langle x, y \mid x^{-1}yx = y^l \rangle$. Arch. Math. 62 (1994), 1–11.
- [8] Carl Droms, Jacques Lewin, and Herman Servatius. The length of elements in free solvable groups. Proc. of the Amer. Math. Soc. 119, no. 1 (1993), 27–33.
- [9] R. Z. Goldstein and E. C. Turner. Fixed subgroups of homomorphisms of free groups. Bull. London Math. Soc. 18 (1986), 468–470.
- [10] K. W. Gruenberg. Residual properties of infinite soluble groups. Proc. London Math. Soc. 7, no. 3 (1957), 29–62.
- [11] V. S. Guba. On the properties of the Cayley graph of Richard Thompson’s group F . Int. J. Alg. Comp. 14, no. 5-6 (2004), 677–702.
- [12] A. J. Guttmann and A. R. Conway. Square lattice self-avoiding walks and polygons. Annals of Combinatorics 5 (2001), 319–345.
- [13] R. Lyndon and P. Schupp. Combinatorial group theory. Springer-Verlag, 1977.
- [14] W. Parry. Growth series of some wreath products. Trans. Amer. Math. Soc. 331, no. 2 (1992), 751–759.
- [15] J.-P. Serre. Trees. Springer-Verlag, 1980.
- [16] A. L. Shmel’kin. Wreath products and varieties of groups (Russian). Izv. Akad. Nauk SSSR Ser. Mat. 29 (1965), 149–170.
- [17] J. Stallings. Group theory and three-dimensional manifolds. Yale Math. Monograph 4. New Haven: Yale Univ. Press, 1971.

VOLOGDA STATE PEDAGOGICAL UNIVERSITY, 6 S. ORLOV STREET, VOLOGDA
160600 RUSSIA

E-mail address: guba@uni-vologda.ac.ru